

Timing and causes of mid-Holocene mammoth extinction on St. Paul Island, Alaska

Russell W. Graham^{a,1}, Soumaya Belmecheri^{a,b}, Kyungcheol Choy^c, Brendan J. Culleton^d, Lauren J. Davies^e, Duane Froese^e, Peter D. Heintzman^f, Carrie Hritz^g, Joshua D. Kapp^f, Lee A. Newsom^{h,i}, Ruth Rawcliffe^c, Émilie Saulnier-Talbot^c, Beth Shapiro^{f,j}, Yue Wang^k, John W. Williams^{k,l}, and Matthew J. Wooller^{c,m}

^aDepartment of Geosciences, College of Earth and Mineral Sciences, The Pennsylvania State University, University Park, PA 16802; ^bLaboratory of Tree Ring Research, University of Arizona, Tucson, AZ 85721; ^cAlaska Stable Isotope Facility, Water and Environmental Research Center, Institute of Northern Engineering, University of Alaska Fairbanks, Fairbanks, AK 99775; ^dHuman Paleoeology and Isotope Geochemistry Laboratory, Department of Anthropology, The Pennsylvania State University, University Park, PA 16802; ^eDepartment of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E3; ^fDepartment of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA 95064; ^gAAAS Science and Technology Policy Fellow, National Science Foundation, Arlington, VA 22230; ^hDepartment of Anthropology, The Pennsylvania State University, University Park, PA 16802; ⁱPenn State Institutes of Energy and the Environment, The Pennsylvania State University, University Park, PA 16802; ^jUC Santa Cruz Genomics Institute, University of California, Santa Cruz, CA 95064; ^kDepartment of Geography, University of Wisconsin–Madison, Madison, WI 53706; ^lCenter for Climatic Research, University of Wisconsin–Madison, Madison, WI 53706; and ^mSchool of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks, AK 99775

Edited by Adrian M. Lister, The Natural History Museum, London, United Kingdom, and accepted by Editorial Board Member David Jablonski June 10, 2016 (received for review March 25, 2016)

Relict woolly mammoth (*Mammuthus primigenius*) populations survived on several small Beringian islands for thousands of years after mainland populations went extinct. Here we present multiproxy paleoenvironmental records to investigate the timing, causes, and consequences of mammoth disappearance from St. Paul Island, Alaska. Five independent indicators of extinction show that mammoths survived on St. Paul until $5,600 \pm 100$ y ago. Vegetation composition remained stable during the extinction window, and there is no evidence of human presence on the island before 1787 CE, suggesting that these factors were not extinction drivers. Instead, the extinction coincided with declining freshwater resources and drier climates between 7,850 and 5,600 y ago, as inferred from sedimentary magnetic susceptibility, oxygen isotopes, and diatom and cladoceran assemblages in a sediment core from a freshwater lake on the island, and stable nitrogen isotopes from mammoth remains. Contrary to other extinction models for the St. Paul mammoth population, this evidence indicates that this mammoth population died out because of the synergistic effects of shrinking island area and freshwater scarcity caused by rising sea levels and regional climate change. Degradation of water quality by intensified mammoth activity around the lake likely exacerbated the situation. The St. Paul mammoth demise is now one of the best-dated prehistoric extinctions, highlighting freshwater limitation as an overlooked extinction driver and underscoring the vulnerability of small island populations to environmental change, even in the absence of human influence.

mammoth | extinction | Holocene | St. Paul Island | ancient DNA

During the wave of late Quaternary extinctions, nearly two-thirds of megafaunal genera disappeared worldwide (1, 2). Proposed extinction drivers include environmental change, human impacts, or some combination, with the relative contributions of these varying by species and region (2). Woolly mammoths, an iconic Ice Age species, vanished from mainland Asia and North America between 14,000 and 13,200 y ago, with some mainland populations perhaps persisting until approximately 10,500 y ago (1, 3, 4); however, relict populations survived on two newly formed Beringian islands into the middle Holocene (4–7) (Fig. 1A).

Wrangel Island is relatively large (7,600 km²), and the late survival of mammoths until 4,020 y ago (5) is consistent with a retreat to refugia in northeastern Siberia as temperatures rose (6). Conversely, the remarkable persistence of mammoths on the much smaller St. Paul Island (110 km²) until the mid-Holocene suggests that isolated megafaunal populations can survive thousands of years after habitat fragmentation. Because there is no evidence of humans on St. Paul Island before the arrival of Russian whalers in 1787 CE (7), the island affords a unique opportunity to study the

processes governing the persistence and extinction of small megafaunal populations in the absence of humans.

St. Paul Island is a remnant of the Bering Land Bridge that became isolated between 14,700 and 13,500 y ago due to sea level rise during the last deglaciation. The island rapidly shrank in area until 9,000 y ago, and then slowly shrank until 6,000 y ago (Fig. 1B). Today, St. Paul is highly isolated (>450 km from Alaska and Aleutians) and is characterized by low relief (maximum elevation 203 m above sea level), a few freshwater lakes, no springs or streams, no permafrost, and moderately productive moss-herbaceous tundra vegetation (8, 9). Apart from a small

Significance

St. Paul Island, Alaska, is famous for its late-surviving population of woolly mammoth. The puzzle of mid-Holocene extinction is solved via multiple independent paleoenvironmental proxies that tightly constrain the timing of extinction to $5,600 \pm 100$ y ago and strongly point to the effects of sea-level rise and drier climates on freshwater scarcity as the primary extinction driver. Likely ecosystem effects of the mega-herbivore extinction include reduced rates of watershed erosion by elimination of crowding around water holes and a vegetation shift toward increased abundances of herbaceous taxa. Freshwater availability may be an underappreciated driver of island extinction. This study reinforces 21st-century concerns about the vulnerability of island populations, including humans, to future warming, freshwater availability, and sea level rise.

Author contributions: Proxy analyses were conducted by R.W.G. (mammoth paleobiology), S.B. (¹⁴C dates and age model), K.C. (stable isotopes and geochemistry), B.J.C. (¹⁴C dates and age model), L.J.D., D.F. (tephra), P.D.H., J.D.K. (sedaDNA), L.A.N. (plant macrofossils), R.R. (cladocerans), É.S.T. (diatoms), B.S. (sedaDNA), Y.W., J.W.W. (pollen and spores), and M.J.W. (stable isotopes and geochemistry); K.C., L.J.D., D.F., P.D.H., L.A.N., B.S., Y.W., J.W.W., and M.J.W. conducted fieldwork, led by R.W.G. and S.B.; K.C. and M.J.W. led paleoenvironmental interpretations; Y.W. conducted statistical analyses; S.B. created core stratigraphy, led sampling strategy, and coordinated data acquisition; R.W.G. coordinated project; C.H. built GIS model of St. Paul Island area; and all authors contributed to manuscript writing, led by R.W.G., P.D.H., and J.W.W.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. A.M.L. is a guest editor invited by the Editorial Board.

Freely available online through the PNAS open access option.

Data deposition: Data have been deposited at the Neotoma Paleocological Database (www.neotomadb.org).

¹To whom correspondence should be addressed. Email: rgraham@ems.psu.edu.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1604903113/-DCSupplemental.

decreased inputs of terrestrial sediments, likely due to reduced rates of watershed erosion after the extinction. Taken together, these proxies strongly indicate lake shallowing, enhanced evaporation, increased turbidity, and increased solute concentrations at Lake Hill during the mid-Holocene extinction window of mammoth on St. Paul Island.

Freshwater availability on oceanic islands is an underappreciated, but important, driver of vertebrate mortality. On Mauritius, a mass mortality event 4,150 y ago coincided with a severe drought that concentrated animals near increasingly toxic and saline lakes (20). Modern elephants (*Loxodonta* and *Elephas*), the closest living equivalents to mammoths, consume 70–200 L/day (21), and their survival depends on water availability (22). Elephants prefer clear water and will dig “wells” near sediment-choked water sources to filter and clarify drinking water (23). St. Paul mammoths may have needed even more water than modern elephants for evaporative cooling during the Holocene interglacial (24), because many of the morphological and physiological adaptations for *M. primigenius* were for retaining heat (25, 26). As island size decreased, freshwater resources on St. Paul became more limited. Surface availability of fresh water is presently—and was probably at the time of extinction—restricted to shallow groundwater perched on a seawater base and several freshwater coastal lagoons and three closed-basin crater lakes (27). Of these, the freshwater lake at Lake Hill is currently the deepest (1.3 m) (9) and thus was the primary (or only) freshwater source for mammoths during dry periods (20). Thus, freshwater availability on St. Paul is highly limited, and diminished freshwater availability is a plausible agent of extinction on St. Paul.

Several previously hypothesized extinction drivers for St. Paul can be discarded. Human colonization during the mid-Holocene is unlikely, given the extreme isolation of St. Paul in the Bering Sea and the lack of any archaeological record (7). Polar bear predation is unlikely, because all St. Paul polar bear remains postdate the mammoth extinction by >1,000 y (11). Volcanic activity (11, 28) is unlikely, given the absence of visible tephra in Lake Hill sediments during the extinction window. Extinction due to increasing winter snowpack (12, 29) is inconsistent with the increase in $\delta^{18}\text{O}$ values at Lake Hill during the Holocene (Fig. 3). Extinction due to vegetation shifts (8) is unlikely, given the stable vegetation composition on St. Paul during the Holocene (Fig. 3), although declining pollen accumulation rates in the early Holocene might indicate declining tundra productivity.

Reduction in island size per se (8) is not a direct driver, because the modern size of the island (110 km²) was reached by 9,000 y ago, more than 3,000 y before extinction. However, decreasing island size may have indirectly caused extinction, by reducing mammoth population sizes and reducing the number and volume of freshwater resources. If so, the St. Paul mammoth extinction may be a natural example of extinction debt (30), and suggests that the delay between habitat loss and final extinction can last for millennia.

Causes of Freshwater Scarcity. The causes of reduced freshwater availability on St. Paul are less certain, with likely factors including sea level rise, climate change, and resource overuse. Rising sea levels gradually decreased land area and availability of freshwater resources (Fig. 1), but a freshwater crisis might have been triggered by mid-Holocene regional climate change and variability. The Lake Hill findings and $\delta^{15}\text{N}$ values from mammoth bones indicate increased aridity between 8,000 and 5,300 y ago on St. Paul (Fig. 3). Other lake records from mainland Alaska show decreased effective moisture and increased hydrological variability between 6,800 and 5,000 y ago (31–34), suggesting that water scarcity of St. Paul might have been caused by regionally enhanced hydrological variability during the mid-Holocene.

Mammoth activity around Lake Hill probably contributed to the degradation of freshwater quality. As ecosystem engineers,

elephants strongly modify their local environments, especially at water holes and other areas of heavy use (35). As plant cover is destroyed, environmental deterioration rapidly radiates outward (12). The abrupt decrease in magnetic susceptibility, shift from terrigenous to lacustrine sediments (Fig. 3), disappearance of pelagic cladocerans, near-disappearance of planktonic diatoms, and presence of tychoplanktonic diatoms all coincide with the time of mammoth extinction. Thus, mammoth activity likely denuded the lake margin, accelerated rates of erosion, and contributed to lake infilling and decreased water quality. The increase in magnetic susceptibility before extinction (Fig. 3) is consistent with intensified water use and rates of watershed erosion, perhaps as other water sources were depleted.

Conclusions and Implications. Multiple reinforcing lines of evidence indicate that reduced freshwater availability triggered the extinction of St. Paul mammoths at $5,600 \pm 100$ y ago. Sea level rise and declining island area set the stage for extinction (8) by reducing island carrying capacity, decreasing population size, and increasing mammoth dependency on diminishing freshwater resources, leaving mammoths vulnerable to environmental fluctuations. Regional climate variability and aridification then may have applied the coup de grâce, possibly exacerbated by resource degradation by the mammoths. Thus, the St. Paul history demonstrates that in the absence of human pressures, small megafaunal populations can persist for thousands of years, while highlighting the susceptibility of island populations to environmental fluctuations, including freshwater availability, particularly during periods of rising sea level. Island biogeographic theory predicts that rates of extinction should be elevated on small islands with limited resources; these findings support that theory while pointing to freshwater scarcity as an underappreciated driver. Our results suggest that as sea levels rise and hydroclimatic variability increases over the coming centuries due to anthropogenic warming (36), freshwater scarcity may become an increasingly critical limiting resource for island vertebrate populations.

Methods

Specific details of all methods are provided in *SI Appendix, SI Text*. Island size through time was reconstructed using a GIS bathymetric map of the St. Paul Island area and the sea level rise curve from the Bering Strait (37). Three overlapping cores were collected at the primary Lake Hill core in 2013 CE (57.17809N, 170.24828W) and used to construct a composite core using clearly identifiable marker layers, such as visible tephra and lithologic transitions, in the overlapping sections (*SI Appendix, Fig. S2*). Samples for all paleoecological and paleoenvironmental proxies were taken at National Lacustrine Core Facility (LacCore) mostly at identical depths, to eliminate problems with proxy correlation. A Bayesian age model for the upper 740 cm of the composite core (*SI Appendix, Fig. S3*) was constructed using OxCal from accelerator mass spectrometry (AMS) radiocarbon (¹⁴C) dating of six terrestrial macrobotanical remains and a tephra layer representing the Aniakhak CFE II tephra, with a known age of $3,595 \pm 4$ cal yr BP (*SI Appendix, Table S3*). Tephra extraction was done following standard methods (38). Radiocarbon dating and isotopic analyses of mammoth teeth were derived from collagen extracted and purified following the modified Longin method with ultrafiltration (39). All AMS ¹⁴C results were corrected for isotopic fractionation according to standard conventions (40). Conventional ¹⁴C ages were calibrated with OxCal 4.2.3 (41) using the IntCal13 curve (42).

SedaDNA from 38 samples throughout the upper 740 cm of the core was extracted, sequenced, and bioinformatically processed at the Paleogenomics Laboratory, University of California Santa Cruz (*SI Appendix, Fig. S5 and Table S5*). Sequencing data are available from the National Center for Biotechnology Information's Short Read Archive (BioProject accession no. PRJNA320875). Extraction and processing of spores and pollen followed a modified version of the LacCore protocol (43). Plant macrofossil samples were taken every 10 cm, but shifted to finer resolution (sampling every 2–5 cm) for the lower portions of the core. All macrofossil samples were disaggregated in a glass Petri dish with distilled water and were manually collected. Samples were not sieved.

Cladocera, diatom, and stable isotopic analyses were all conducted at the Alaska Stable Isotope Facility, University of Alaska Fairbanks. Preparation of

diatom samples was done following a modified previously published protocol (44). Cladocera were processed with an adapted version of a method described previously (45). The $\delta^{18}\text{O}$ values of chironomid head capsules and other aquatic insect chitin were processed and analyzed following previously published protocols (46, 47). Bulk sediment samples were collected from the core at 16-cm intervals for isotopic and other geochemical analyses and processed following previously published protocols (48).

ACKNOWLEDGMENTS. We thank all of the inhabitants of St. Paul Island and in particular Jason Bourdukofski, Bill Briggs, Aquilina Lestenkof, Gary Stanley, Mac Mandregan, Phillip Zavadi, Brenda Jones, Connie Newman, and the National Oceanic and Atmospheric Administration for their

hospitality, assistance with site access, and field logistics. Simeon Swetozof, Laura Lestenkof, Nick Kozloff, Rob Owens, and Logan Tetov kindly provided specimens of woolly mammoth fossils from personal collections for radiocarbon dating. Voucher specimens have been preserved in the Earth and Mineral Sciences Museum collections at the Pennsylvania State University. The Limnological Research Center provided facilities for core splitting, imaging, description, and sampling. Elle Palkopoulou and Love Dalén provided prepublication access to the Wrangel Island woolly mammoth genome. We thank Nick Holschuh for assistance with figure drafting, Nancy Bigelow and Greg Smith for assistance with fieldwork, and two anonymous reviewers for their careful review and comments. This work was supported by National Science Foundation Grants PLR-1204233, PLR-1203772, and PLR-1203990.

- Guthrie RD (2006) New carbon dates link climatic change with human colonization and Pleistocene extinctions. *Nature* 441(7090):207–209.
- Barnosky AD, Koch PL, Feranec RS, Wing SL, Shabel AB (2004) Assessing the causes of late Pleistocene extinctions on the continents. *Science* 306(5693):70–75.
- Haile J, et al. (2009) Ancient DNA reveals late survival of mammoth and horse in interior Alaska. *Proc Natl Acad Sci USA* 106(52):22352–22357.
- Nikoskiy PA, Sulerzhitsky LD, Pitulko VV (2011) Last straw versus Blitzkrieg overkill: Climate-driven changes in the Arctic Siberian mammoth population and the Late Pleistocene extinction problem. *Quat Sci Rev* 30:2309–2328.
- Vartanyan SL, Arslanov KA, Karhu JA, Possnert G, Sulerzhitsky LD (2008) Collection of radiocarbon dates on the mammoths (*Mammuthus primigenius*) and other genera of Wrangel Island, northeast Siberia, Russia. *Quat Res* 70(1):51–59.
- Lorenzen ED, et al. (2011) Species-specific responses of Late Quaternary megafauna to climate and humans. *Nature* 479(7373):359–364.
- Veltre DW, Veltre MJ (1981) A preliminary baseline study of subsistence resource utilization in the Pribilof Islands. Alaska Department of Fish and Game, Division of Subsistence, Technical Paper no. 57. Available at: www.subsistence.adfg.state.ak.us/TechPap/tp057.pdf. Accessed June 22, 2016.
- Guthrie RD (2004) Radiocarbon evidence of mid-Holocene mammoths stranded on an Alaskan Bering Sea island. *Nature* 429(6993):746–749.
- Colinvaux P (1981) Historical ecology in Beringia: The south land bridge coast at St. Paul Island. *Quat Res* 16(1):18–36.
- Hanna GD (1922) The reindeer herds of the Pribilof Islands. *Sci Mon* 15:181–186.
- Veltre DW, Yesner DR, Crossen KJ, Graham RW, Coltrain JB (2008) Patterns of faunal extinction and paleoclimatic change from mid-Holocene mammoth and polar bear remains, Pribilof Islands, Alaska. *Quat Res* 70(1):40–50.
- Bryson RB, Agenbroad LD, McEnaney DeWall K (2010) Paleoclimate modeling and paleoenvironmental interpretations for three instances of island dwelling mammoths. *Quat Int* 217(1–2):6–9.
- Willerslev E, et al. (2003) Diverse plant and animal genetic records from Holocene and Pleistocene sediments. *Science* 300(5620):791–795.
- Baker AG, Bhagwat SA, Willis KJ (2013) Do dung fungal spores make a good proxy for past distribution of large herbivores? *Quat Sci Rev* 62:21–31.
- Rawlence NJ, et al. (2014) Using palaeoenvironmental DNA to reconstruct past environments: Progress and prospects. *J Quat Sci* 29(7):610–626.
- van Geel B, Zazula GD, Schweger CE (2007) Spores of coprophilous fungi from under the Dawson tephra (25,300 ^{14}C years BP), Yukon Territory, northwestern Canada. *Palaeogeogr Palaeoclimatol Palaeoecol* 252:481–485.
- Thienpont JR, et al. (2013) Exploratory hydrocarbon drilling impacts to Arctic lake ecosystems. *PLoS One* 8(11):e78875.
- Stevens RE, Hedges REM (2004) Carbon and nitrogen stable isotope analysis of northwest European horse bone and tooth collagen, 40,000 BP–present: Palaeoclimatic interpretations. *Quat Sci Rev* 23(7–8):977–991.
- Newsome SD, Miller GH, Magee JW, Fogel ML (2011) Quaternary record of aridity and mean annual precipitation based on $\delta^{15}\text{N}$ in ratite and dromornithid eggshells from Lake Eyre, Australia. *Oecologia* 167(4):1151–1162.
- de Boer EJ, et al. (2015) A deadly cocktail: How a drought around 4200 cal yr BP caused mass mortality events at the infamous “dodo swamp” in Mauritius. *Holocene* 25(5):758–771.
- Benedict FG (1936) *The Physiology of the Elephant* (The Carnegie Institution of Washington, Washington, DC).
- Western D (1975) Water availability and its influence on the structure and dynamics of a savanna large mammal community. *E Afr Wildl J* 13:265–286.
- Ramey EM, Ramey RR, Brown LM, Kelley ST (2013) Desert-dwelling African elephants (*Loxodonta africana*) in Namibia dig wells to purify drinking water. *Pachyderm* 53:66–72.
- Dunkin RC, Wilson D, Way N, Johnson K, Williams TM (2013) Climate influences thermal balance and water use in African and Asian elephants: Physiology can predict drivers of elephant distribution. *J Exp Biol* 216(Pt 15):2939–2952.
- Kubiak H (1982) Morphological characters of the mammoth: an adaptation to the arctic-steppe environment. *Paleoecology of Beringia*, eds Hopkins DM, Matthews JV, Schweger CE (Academic, New York), pp 281–289.
- Tridico SR, Rigby P, Kirkbride KP, Haile J, Bunce M (2014) Megafaunal split ends: Microscopical characterisation of hair structure and function in extinct woolly mammoth and woolly rhino. *Quat Sci Rev* 83:68–75.
- Feulner AJ (1980) Water resources reconnaissance of the southeastern part of St. Paul Island. USGS Water-Resources Investigations Report 80-61 (US Geological Survey, Reston, VA).
- Clegg BF, Hu FS (2010) An oxygen-isotope record of Holocene climate change in the south-central Brooks Range, Alaska. *Quat Sci Rev* 29(7–8):928–939.
- Finney BP, Bigelow NH, Barber VA, Edwards ME (2012) Holocene climate change and carbon cycling in a groundwater-fed, boreal forest lake: Dune Lake, Alaska. *J Paleolimnol* 48:43–54.
- Kuusaaari M, et al. (2009) Extinction debt: A challenge for biodiversity conservation. *Trends Ecol Evol* 24(10):564–571.
- Jones MC, Wooller M, Peteet DM (2014) A deglacial and Holocene record of climate variability in south-central Alaska from stable oxygen isotopes and plant macrofossils in peat. *Quat Sci Rev* 87:1–11.
- Laws RM (1970) Elephants as agents of habitat and landscape change in East Africa. *Oikos* 21:1–15.
- Van Wyk P, Fairall N (1969) The influence of the African elephant on the vegetation of the Kruger National Park. *Koedoe* 12:57–89.
- Mayewski PA, et al. (2004) Holocene climate variability. *Quat Res* 62(3):243–255.
- Feeley TC, Winer GS (2009) Volcano hazards and potential risks on St. Paul Island, Pribilof Islands, Bering Sea, Alaska. *J Volcanol Geotherm Res* 182(1–2):57–66.
- Clark PU, et al. (2016) Consequences of twenty-first-century policy for multi-millennial climate and sea-level change. *Nat Clim Change* 6:360–369.
- Hu A, et al. (2010) Influence of Bering Strait flow and North Atlantic circulation on glacial sea-level changes. *Nat Geosci* 3(2):118–121.
- Blockley SPE, et al. (2005) A new and less destructive laboratory procedure for the physical separation of distal glass tephra shards from sediments. *Quat Sci Rev* 24(16–17):1952–1960.
- Brown TA, Nelson DE, Vogel JS, Southon JR (1988) Improved collagen extraction by modified Longin method. *Radiocarbon* 30(2):171–177.
- Stuiver M, Polach HA (1977) Discussion: Reporting of ^{14}C data. *Radiocarbon* 19(3):355–363.
- Bronk RC (2009) Bayesian analysis of radiocarbon dates. *Radiocarbon* 51:337–360.
- Reimer PJ, et al. (2013) Intcal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon* 55(4):1869–1887.
- Faegri K, Kaland PE, Krzywinski K (1989) *Textbook of Pollen Analysis* (Wiley, New York), 4th Ed.
- Battarbee RW, et al. (2001) Diatoms. Tracking Environmental Change Using Lake Sediments, Vol. 3: Terrestrial, Algal and Siliceous Indicators, eds Smol JP, Birks HJB, Last WM (Kluwer, Dordrecht), pp 155–202.
- Korhola A, Rautio M (2001) Cladocera and other branchiopod crustaceans. Tracking Environmental Change Using Lake Sediments, Vol. 4: Zoological Indicators, eds Smol JP, Birks HJB, Last WM (Kluwer, Dordrecht), pp 5–41.
- Wooller MJ, et al. (2004) Quantitative paleotemperature estimates from $\delta^{18}\text{O}$ of chironomid head capsules preserved in arctic lake sediments. *J Paleolimnol* 31(3):267–274.
- Wooller MJ, et al. (2012) Reconstruction of past methane availability in an Arctic Alaska wetland indicates climate influenced methane release during the past similar to 12,000 years. *J Paleolimnol* 48(1):27–42.
- Wooller MJ, et al. (2012) An ~11,200-year paleolimnological perspective for emerging archaeological findings at Quartz Lake, Alaska. *J Paleolimnol* 48(1):83–99.